



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### On the magnitude of a finite dimensional algebra

**Citation for published version:**

Chuang, J, King, A & Leinster, T 2016, 'On the magnitude of a finite dimensional algebra', *Theory and Applications of Categories*. <<http://www.tac.mta.ca/tac/volumes/31/3/31-03abs.html>>

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Published In:**

Theory and Applications of Categories

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



# ON THE MAGNITUDE OF A FINITE DIMENSIONAL ALGEBRA

JOSEPH CHUANG, ALASTAIR KING AND TOM LEINSTER

**ABSTRACT.** There is a general notion of the magnitude of an enriched category, defined subject to hypotheses. In topological and geometric contexts, magnitude is already known to be closely related to classical invariants such as Euler characteristic and dimension. Here we establish its significance in an algebraic context. Specifically, in the representation theory of an associative algebra  $A$ , a central role is played by the indecomposable projective  $A$ -modules, which form a category enriched in vector spaces. We show that the magnitude of that category is a known homological invariant of the algebra: writing  $\chi_A$  for the Euler form of  $A$  and  $S$  for the direct sum of the simple  $A$ -modules, it is  $\chi_A(S, S)$ .

## 1. Introduction

This paper is part of a large programme to define and investigate cardinality-like invariants of mathematical objects. Given a monoidal category  $\mathbf{V}$  together with a notion of the ‘size’  $|X|$  of each object  $X$  of  $\mathbf{V}$ , there arises automatically a notion of the ‘size’ or ‘magnitude’ of each  $\mathbf{V}$ -category (subject to conditions). Here we apply this general method in the context of associative algebras.

More specifically, for any finite-dimensional algebra  $A$ , the category  $\mathbf{IP}(A)$  of indecomposable projective  $A$ -modules plays a central role (discussed below) in the theory of representations of  $A$ . This category is enriched in finite-dimensional vector spaces, and, taking dimension as the base notion of size, we can then consider the magnitude of  $\mathbf{IP}(A)$ . We show that this is a known homological invariant of the original algebra  $A$ .

Little algebra will be assumed on the reader’s part; all the necessary background is provided in Section 2.

The general definition of magnitude is as follows [9, §1.3]. Let  $\mathbf{V}$  be a monoidal category equipped with a function  $|\cdot|$  on its set of objects (taking values in a semiring, say). Let  $\mathbf{A}$  be a  $\mathbf{V}$ -category with finitely many objects. Denote by  $Z_{\mathbf{A}} = (Z_{ab})$  the square matrix whose rows and columns are indexed by the objects of  $\mathbf{A}$ , and with entries

$$Z_{ab} = |\mathbf{A}(a, b)| \tag{1}$$

( $a, b \in \mathbf{A}$ ). If  $Z_{\mathbf{A}}$  is invertible, the **magnitude**  $|\mathbf{A}|$  of  $\mathbf{A}$  is defined to be the sum of all the entries of  $Z_{\mathbf{A}}^{-1}$ .

---

Received by the editors 2015-06-11 and, in final form, 2016-01-06.

Transmitted by Ross Street. Published on 2016-01-07.

2010 Mathematics Subject Classification: 18G99 (primary), 16D40, 16D60, 16G99, 18E05, 18G15.

Key words and phrases: algebra, magnitude, indecomposable projective, simple module, Cartan matrix, Euler form, Cartan determinant conjecture.

© Joseph Chuang, Alastair King and Tom Leinster, 2015. Permission to copy for private use granted.

Since  $Z_{\mathbf{A}}$  need not be invertible, magnitude is not defined for every  $\mathbf{A}$ . But where magnitude *is* defined, we may harmlessly extend the definition by equivalence, setting  $|\mathbf{A}| = |\mathbf{B}|$  whenever  $\mathbf{A}$  and  $\mathbf{B}$  are equivalent  $\mathbf{V}$ -categories such that  $\mathbf{B}$  has finitely many objects and  $Z_{\mathbf{B}}$  is invertible. (There is no problem of consistency, since if  $\mathbf{A}$  and  $\mathbf{B}$  are equivalent and both  $Z_{\mathbf{A}}$  and  $Z_{\mathbf{B}}$  are invertible then both  $\mathbf{A}$  and  $\mathbf{B}$  are skeletal—that is, isomorphic objects are equal—and so  $\mathbf{A}$  and  $\mathbf{B}$  are isomorphic.)

Unmotivated as this definition may seem, multiple theorems attest that magnitude is the canonical notion of the size of an enriched category. For example, take  $\mathbf{V}$  to be the category of finite sets and  $|X|$  to be the cardinality of a finite set  $X$ . Then we obtain a notion of the magnitude of a finite category. In this context, magnitude is also called Euler characteristic [8], for the following reason. Recall that every small category  $\mathbf{A}$  gives rise to a topological space  $B\mathbf{A}$ , its classifying space or geometric realisation. Proposition 2.11 of [8] states that under finiteness hypotheses,

$$|\mathbf{A}| = \chi(B\mathbf{A}). \quad (2)$$

Thus, the Euler characteristic of a category has a similar status to group (co)homology: it is defined combinatorially, but agrees with the topological notion when one passes to the classifying space.

For another example, let  $\mathbf{V}$  be the ordered set  $([0, \infty], \geq)$  with addition as the monoidal structure, so that metric spaces can be viewed as  $\mathbf{V}$ -categories [6]. For  $x \in [0, \infty]$ , put  $|x| = e^{-x}$ . (The virtue of this choice is that  $|x \otimes y| = |x| |y|$ .) Then we obtain a notion of the magnitude of a finite metric space. This extends naturally to a large class of compact metric spaces [9, 11, 12]. The magnitude of a compact subset of  $\mathbb{R}^n$  is always well-defined, and is closely related to classical quantities of geometric measure. For example, a theorem of Meckes [12, Corollary 7.4] shows that Minkowski dimension can be recovered from magnitude, and conjectures of Leinster and Willerton [10] state that magnitude also determines invariants such as volume and surface area.

Here we study the case where  $\mathbf{V}$  is the category of finite-dimensional vector spaces and  $|X| = \dim X$ . We then obtain a notion of the magnitude of a linear (that is,  $\mathbf{V}$ -enriched) category. Our main theorem is this:

**1.1. THEOREM.** *Let  $A$  be an algebra of finite dimension and finite global dimension over an algebraically closed field. Write  $\mathbf{IP}(A)$  for the linear category of indecomposable projective  $A$ -modules,  $(S_i)_{i \in I}$  for representatives of the isomorphism classes of simple  $A$ -modules, and  $S = \bigoplus_{i \in I} S_i$ . Then*

$$|\mathbf{IP}(A)| = \sum_{n=0}^{\infty} (-1)^n \dim \operatorname{Ext}_A^n(S, S). \quad (3)$$

We now explain the context of this result; background can be found in the next section.

Any associative algebra  $A$  gives rise to several linear categories, including the category of all  $A$ -modules and the one-object category corresponding to  $A$  itself (which trivially has magnitude  $1/\dim A$ ). But it also gives rise to the category  $\mathbf{IP}(A)$  of indecomposable

projective  $A$ -modules, whose main significance is that its representation theory is the same as that of  $A$ :

$$\begin{aligned} A\text{-}\mathbf{Mod} &\simeq [\mathbf{IP}(A)^{\mathrm{op}}, \mathbf{Vect}] \\ M &\mapsto \mathrm{Hom}_A(-, M) \end{aligned} \tag{4}$$

where the right-hand side is the category of contravariant linear functors from  $\mathbf{IP}(A)$  to vector spaces. In other words,  $\mathbf{IP}(A)^{\mathrm{op}}$  and the one-object linear category  $A$  are Morita equivalent.

The Krull–Schmidt theorem states that every finitely generated  $A$ -module can be expressed as a direct sum of indecomposable modules, in an essentially unique way. It implies that the  $A$ -module  $A$  is a direct sum of indecomposable *projective* modules, and that, moreover, every indecomposable projective appears at least once in this sum. Thus, the indecomposable projectives are the ‘atoms’ of  $A$ , in the sense of being its constituent parts.

This explains the equivalence (4). The absolute colimits in linear categories are the finite direct sums and idempotent splittings (that is, direct summands). Every finitely generated projective module is a direct sum of indecomposable projectives, so the category of finitely generated projectives is the Cauchy completion of  $\mathbf{IP}(A)$ . On the other hand, every finitely generated projective is a direct summand of a direct sum of copies of the  $A$ -module  $A$ , so the category of finitely generated projectives is also the Cauchy completion of the one-object category  $A^{\mathrm{op}}$ . Hence  $\mathbf{IP}(A)$  and  $A^{\mathrm{op}}$  have the same Cauchy completion, and are therefore Morita equivalent.

The simple modules, too, can be thought of as ‘atomic’ in a different sense. A simple module need not be indecomposable projective, nor vice versa. However, the two conditions are closely related: as recounted in Section 2, there is a canonical bijection between the isomorphism classes of simple modules and the isomorphism classes of indecomposable projectives.

The condition that  $A$  has finite global dimension guarantees that the sum in (3) has only finitely many nonzero terms. The condition that  $A$  has finite dimension guarantees that the linear category  $\mathbf{IP}(A)$  is equivalent to one with finitely many objects and finite-dimensional hom-spaces, as we shall see. This is a necessary condition in order for the magnitude of  $\mathbf{IP}(A)$  to be defined. It is not a sufficient condition, but part of the statement of Theorem 1.1 is that  $|\mathbf{IP}(A)|$  is defined.

Theorem 1.1 was first noted by Catharina Stroppel under the additional hypothesis that  $A$  is a Koszul algebra (personal communication, 2009). We observe here that the Koszul assumption is unnecessary.

## 2. Algebraic background

Here we assemble all the facts that we will need in order to state and prove the main theorem. General references for this section are [13, Chapter I] and [2, Chapter 1].

Throughout this note,  $K$  denotes a field and  $A$  a finite-dimensional  $K$ -algebra (unital, but not necessarily commutative). ‘Module’ will mean left  $A$ -module. Since  $A$  is finite-

dimensional, a module is finitely generated over  $A$  if and only if it is finite-dimensional over  $K$ .

**SIMPLE AND INDECOMPOSABLE PROJECTIVE MODULES.** Details for this part can be found in [7], as well as in the general references above.

A nonzero module is **simple** if it has no nontrivial submodule, and **indecomposable** if it has no nontrivial direct summand. There is a canonical bijection between the isomorphism classes of simple modules  $S$  and the isomorphism classes of indecomposable projective modules  $P$ , with  $S$  corresponding to  $P$  if and only if  $S$  is a quotient of  $P$ . (It is not an equivalence of categories.)

Choose representatives  $(S_i)_{i \in I}$  of the isomorphism classes of simple modules and  $(P_i)_{i \in I}$  of the isomorphism classes of indecomposable projective modules, with  $S_i$  a quotient of  $P_i$ .

Modules of both types are finitely generated (indeed, cyclic), so each vector space  $\text{Hom}_A(P_i, P_j)$  is finite-dimensional. Moreover, one can use either the Jordan–Hölder theorem or the Krull–Schmidt theorem to show that  $I$  is finite. Denote by  $\mathbf{IP}(A)$  the category of indecomposable projective  $A$ -modules and all homomorphisms between them, which is a  $K$ -linear category. Then  $\mathbf{IP}(A)$  has finite-dimensional hom-spaces and only finitely many isomorphism classes of objects.

We have  $\text{Hom}_A(P_i, S_j) = 0$  when  $i \neq j$ , since any homomorphism into a simple module is zero or surjective. It can be shown that  $\text{Hom}_A(P_i, S_i) \cong \text{End}_A(S_i)$  as vector spaces. This is a skew field, isomorphic to  $K$  if  $K$  is algebraically closed.

**HOMOLOGICAL ALGEBRA.** For each  $n \geq 0$ , there is a functor

$$\text{Ext}_A^n: A\text{-}\mathbf{Mod}^{\text{op}} \times A\text{-}\mathbf{Mod} \rightarrow \mathbf{Vect}. \quad (5)$$

One can characterise  $\text{Ext}_A^n(X, -)$  as the  $n$ th right derived functor of  $\text{Hom}_A(X, -)$ , and  $\text{Ext}_A^n(-, Y)$  as the  $n$ th right derived functor of  $\text{Hom}_A(-, Y)$ , but we will only need the following consequences of these characterisations.

First,  $\text{Ext}_A^0 = \text{Hom}_A$ . Second, if  $P$  is projective then  $\text{Ext}_A^n(P, -) = 0$  for all  $n > 0$ . Third,  $\text{Ext}_A^n$  preserves finite direct sums in each argument. Fourth,  $\text{Ext}_A^n(X, Y)$  is finite-dimensional if both  $X$  and  $Y$  are. Finally, given any  $A$ -module  $V$  and short exact sequence

$$0 \rightarrow W \rightarrow X \rightarrow Y \rightarrow 0, \quad (6)$$

there is an induced long exact sequence

$$\begin{aligned} 0 \rightarrow \text{Ext}_A^0(V, W) \rightarrow \text{Ext}_A^0(V, X) \rightarrow \text{Ext}_A^0(V, Y) \\ \rightarrow \text{Ext}_A^1(V, W) \rightarrow \text{Ext}_A^1(V, X) \rightarrow \cdots, \end{aligned} \quad (7)$$

and dually a long exact sequence  $0 \rightarrow \text{Ext}_A^0(Y, V) \rightarrow \cdots$ .

Assume henceforth that  $A$  has **finite global dimension** [14, Chapter 4]. This means that there exists  $N \in \mathbb{N}$  such that every  $A$ -module  $X$  has a projective resolution of the form

$$0 \rightarrow Q_N \rightarrow \cdots \rightarrow Q_1 \rightarrow X \rightarrow 0. \quad (8)$$

When  $X$  is finite-dimensional, the projective modules  $Q_i$  can be chosen to be finite-dimensional too.

A condition equivalent to finite global dimension is that  $\text{Ext}_A^n = 0$  for all  $n \gg 0$ . For finite-dimensional  $A$ -modules  $X$  and  $Y$ , we may therefore define

$$\chi_A(X, Y) = \sum_{n=0}^{\infty} (-1)^n \dim \text{Ext}_A^n(X, Y) \in \mathbb{Z} \quad (9)$$

(a finite sum). This  $\chi_A$  is the **Euler form** of  $A$ . We have  $\chi_A(\bigoplus_r X_r, -) = \sum_r \chi_A(X_r, -)$  for any finite family  $(X_r)$  of modules, and similarly in the second argument. Moreover, the observations above imply that

$$\chi_A(P_i, P_j) = \dim \text{Hom}_A(P_i, P_j) \quad (10)$$

for all  $i, j \in I$ , and that

$$\chi_A(P_i, S_j) = \begin{cases} \dim \text{End}_A(S_j) & \text{if } i = j, \\ 0 & \text{if } i \neq j. \end{cases} \quad (11)$$

When  $K$  is algebraically closed,  $\chi_A(P_i, S_j)$  is therefore just the Kronecker delta  $\delta_{ij}$ .

**GROTHENDIECK GROUP.** The **Grothendieck group**  $K(A)$  is the abelian group generated by the finite-dimensional  $A$ -modules, subject to the relation  $X = W + Y$  for each short exact sequence (6) of finite-dimensional modules. Writing  $[X]$  for the class of  $X$  in  $K(A)$ , one easily deduces that, more generally,  $\sum_r (-1)^r [X_r] = 0$  for any exact sequence

$$0 \rightarrow X_1 \rightarrow \cdots \rightarrow X_n \rightarrow 0. \quad (12)$$

For example, take a short exact sequence (6) and a finite-dimensional module  $V$ . The resulting long exact sequence (7) has only finitely many nonzero terms (since  $A$  has finite global dimension), so the alternating sum of the dimensions of these terms is 0, giving  $\chi_A(V, X) = \chi_A(V, W) + \chi_A(V, Y)$ . The same holds with the arguments reversed. Thus,  $\chi_A$  defines a  $\mathbb{Z}$ -bilinear map  $K(A) \times K(A) \rightarrow \mathbb{Z}$ .

We now show that  $K(A)$  is free as a  $\mathbb{Z}$ -module, and in fact has two canonical bases.

First, the family  $([S_i])_{i \in I}$  generates the group  $K(A)$ . Indeed, for any finite-dimensional  $A$ -module  $X$ , we may take a composition series

$$0 = X_n < \cdots < X_1 < X_0 = X, \quad (13)$$

and then  $[X] = \sum_{r=1}^n [X_{r-1}/X_r]$ .

Second, the family  $([P_i])_{i \in I}$  generates  $K(A)$ . Given a finite-dimensional  $A$ -module  $X$ , we may take a resolution (8) by finite-dimensional projective modules, and then  $[X] = \sum_{r=1}^N (-1)^{r+1} [Q_r]$ . On the other hand, each  $Q_r$  is a finite direct sum of indecomposable submodules, which are projective since  $Q_r$  is.

Finally, both  $([S_i])$  and  $([P_i])$  *freely* generate the abelian group  $K(A)$ . This follows from (11) and the  $\mathbb{Z}$ -bilinearity of  $\chi_A$ .

### 3. The result

Recall our standing conventions:  $A$  is an algebra of finite dimension and finite global dimension, over a field  $K$  which we now assume to be algebraically closed. We continue to write  $(P_i)_{i \in I}$  for representatives of the isomorphism classes of indecomposable projective  $A$ -modules, and similarly  $(S_i)_{i \in I}$  for the simple modules, with  $S_i$  a quotient of  $P_i$ .

The linear category  $\mathbf{IP}(A)$  of indecomposable projective  $A$ -modules is equivalent to its full subcategory with objects  $P_i$  ( $i \in I$ ). Write  $Z_A = (Z_{ij})_{i,j \in I}$  for the matrix of this finite linear category, so that  $Z_{ij} = \dim \operatorname{Hom}_A(P_i, P_j)$ .

We will derive our main result, Theorem 1.1, from the following basic theorem. (See e.g. [1, Proposition III.3.13(a)] for an essentially equivalent formulation.) It implies, in particular, that the matrix  $Z_A$  is invertible over the integers.

**3.1. THEOREM.** *The inverse of the matrix  $Z_A$  is the ‘Euler matrix’  $E_A = (E_{ij})_{i,j \in I}$ , given by  $E_{ij} = \chi_A(S_j, S_i)$ .*

**PROOF.** Since  $([P_i])_{i \in I}$  and  $([S_i])_{i \in I}$  are both bases for the  $\mathbb{Z}$ -module  $K(A)$ , there is an invertible matrix  $C_A = (C_{ij})_{i,j \in I}$  over  $\mathbb{Z}$  such that, writing  $C_A^{-1} = (\overline{C}_{ij})$ ,

$$[P_j] = \sum_{k \in I} C_{kj} [S_k], \quad (14)$$

$$[S_j] = \sum_{k \in I} \overline{C}_{kj} [P_k] \quad (15)$$

for all  $j \in I$ . Since  $K$  is algebraically closed, equation (11) states that  $\chi_A(P_i, S_j) = \delta_{ij}$ . Applying  $\chi_A(P_i, -)$  to each side of (14) therefore gives  $\chi_A(P_i, P_j) = C_{ij}$ , which by (10) is equivalent to  $Z_{ij} = C_{ij}$ . On the other hand, applying  $\chi_A(-, S_i)$  to each side of (15) gives  $E_{ij} = \overline{C}_{ij}$ . Hence  $Z_A = C_A$  and  $E_A = C_A^{-1}$ . ■

The matrix  $C_A = Z_A$  is known as the **Cartan matrix** of  $A$  ([3], [4, §5], [5]). Explicitly,  $C_{ij}$  is the multiplicity of  $S_i$  as a composition factor of  $P_j$ .

We now deduce Theorem 1.1. By definition,  $|\mathbf{IP}(A)|$  is the sum of the entries of  $Z_A^{-1}$ . Hence by Theorem 3.1 and the  $\mathbb{Z}$ -bilinearity of  $\chi_A$ ,

$$|\mathbf{IP}(A)| = \sum_{i,j \in I} \chi_A(S_j, S_i) = \chi_A\left(\bigoplus_{j \in I} S_j, \bigoplus_{i \in I} S_i\right) = \chi_A(S, S), \quad (16)$$

completing the proof.

**3.2. EXAMPLE.** Let  $Q$  be a finite acyclic quiver (directed graph). Then  $Q$  consists of a finite set  $I$  of vertices together with, for each  $i, j \in I$ , a finite set  $Q(i, j)$  of arrows from  $i$  to  $j$ . The **path algebra**  $A$  of  $Q$  is defined as follows. As a vector space, it is generated by the paths in  $Q$ , including the zero-length path  $e_i$  on each vertex  $i$ . Multiplication is concatenation of paths where that is defined, and zero otherwise. We write multiplication in the same order as composition, so that if  $\alpha$  is a path from  $i$  to  $j$  and  $\beta$  is a path from



$j$  to  $k$  then  $\beta\alpha$  is a path from  $i$  to  $k$ . The identity is  $\sum_{i \in I} e_i$ . That  $Q$  is finite and acyclic guarantees that  $A$  is of finite dimension and finite global dimension.

Path algebras of quivers are very well-understood (e.g. [13, Chapter I]). The simple and indecomposable projective  $A$ -modules are indexed by the vertex-set  $I$ . The indecomposable projective module  $P_i$  corresponding to vertex  $i$  is the submodule of the  $A$ -module  $A$  spanned by the paths beginning at  $i$ . It has a unique maximal submodule  $N_i$ , spanned by the paths of nonzero length beginning at  $i$ , and the corresponding simple module  $S_i = P_i/N_i$  is one-dimensional.

Using the facts listed in Section 2, we can compute the Euler form of  $A$ . For each  $i, j \in I$ , the short exact sequence

$$0 \rightarrow N_i \rightarrow P_i \rightarrow S_i \rightarrow 0 \quad (17)$$

gives rise to a long exact sequence

$$0 \rightarrow \text{Ext}_A^0(S_i, S_j) \rightarrow \text{Ext}_A^0(P_i, S_j) \rightarrow \text{Ext}_A^0(N_i, S_j) \rightarrow \cdots \quad (18)$$

Observing that  $N_i = \bigoplus_{k \in I} P_k^{Q(i,k)}$ , we deduce from (18) that

$$\text{Ext}_A^n(S_i, S_j) = \begin{cases} K^{\delta_{ij}} & \text{if } n = 0, \\ K^{Q(i,j)} & \text{if } n = 1, \\ 0 & \text{if } n \geq 2. \end{cases} \quad (19)$$

Hence, writing  $E = \coprod_{i,j \in Q} Q(i,j)$  for the set of arrows of  $Q$ ,

$$\text{Ext}_A^n(S, S) = \begin{cases} K^{|I|} & \text{if } n = 0, \\ K^{|E|} & \text{if } n = 1, \\ 0 & \text{if } n \geq 2. \end{cases} \quad (20)$$

It follows that  $\chi_A(S, S) = |I| - |E|$ , which is the Euler characteristic (in the elementary sense) of the quiver  $Q$ .

On the other hand, each path from vertex  $j$  to vertex  $i$  induces a homomorphism  $P_i \rightarrow P_j$  by composition, and in fact every homomorphism  $P_i \rightarrow P_j$  is a unique linear combination of homomorphisms of this form. Hence  $Z_{ij}$  is the number of paths from  $j$  to  $i$  in  $Q$ .

So in the case at hand, Theorem 1.1 states that if we take an acyclic quiver  $Q$ , form the matrix whose  $(i, j)$ -entry is the number of paths from  $j$  to  $i$ , invert this matrix, and sum its entries, the result is equal to the Euler characteristic of  $Q$ . This was also shown directly as Proposition 2.10 of [8].

#### 4. Some remarks

ARBITRARY BASE FIELDS. The assumption that the base field is algebraically closed is needed for the simple form of the duality formula,  $\chi_A(P_i, S_j) = \delta_{ij}$ . Otherwise, equa-



tion (11) only gives

$$\chi_A(P_i, S_j) = \begin{cases} d_j & \text{if } i = j, \\ 0 & \text{if } i \neq j, \end{cases} \quad (21)$$

where  $d_j = \dim \text{End}_A(S_j)$ . Then, applying  $\chi_A(P_i, -)$  to (14) yields  $Z_{ij} = d_i C_{ij}$ , while applying  $\chi_A(-, S_i)$  to (15) yields  $E_{ij} = d_i \bar{C}_{ij}$ . Therefore, writing  $Z_A^{-1} = (\bar{Z}_{ij})$ , we get

$$\bar{Z}_{ij} = d_i^{-1} E_{ij} d_j^{-1}, \quad (22)$$

which generalises Theorem 3.1. We can then sum (22) to generalise Theorem 1.1 as follows:

$$|\mathbf{IP}(A)| = \chi_A(\tilde{S}, \tilde{S}), \quad (23)$$

where  $\tilde{S} = \bigoplus_{i \in I} d_i^{-1} S_i$ , which may be regarded as a formal module or we may note that (23) only depends on the class  $[\tilde{S}] = \sum_{i \in I} d_i^{-1} [S_i] \in K(A) \otimes_{\mathbb{Z}} \mathbb{Q}$ .

**THE DETERMINANT OF THE CARTAN MATRIX.** The fact that, when  $A$  has finite global dimension, the Cartan matrix  $C_A$  is invertible over  $\mathbb{Z}$  or, equivalently, is unimodular, i.e.  $\det C_A = \pm 1$ , is an old observation of Eilenberg [4, §5]. On the other hand, the ‘Cartan determinant conjecture’ that, in fact,  $\det C_A = 1$  is still unsolved in general, although it is confirmed in many cases; see [5] for a survey.

An easy example is when  $A$  is (Morita equivalent to) a quotient of the path algebra of an acyclic quiver, in which case  $A$  is necessarily finite dimensional and of finite global dimension. In this case  $C_A = Z_A$  can be made upper triangular with 1s on the diagonal, so it certainly has  $\det C_A = 1$ . As another example, Zacharia [15] showed that the conjecture holds whenever  $A$  has global dimension 2.

It is not hard to give an example of an algebra  $A$  for which  $C_A = Z_A$  is not even invertible over  $\mathbb{Q}$ : e.g. the quiver algebra given by a single  $n$ -cycle, with all paths of length  $n$  set to 0, has  $C_{ij} = Z_{ij} = 1$  for all  $i, j \in I$ . Inevitably, this algebra does not have finite global dimension.

In this example, it is in fact still possible [9, §1] to define the magnitude of  $\mathbf{IP}(A)$ , and indeed  $|\mathbf{IP}(A)| = 1$ . However, it is less clear how one might find a homological interpretation of this.

**ACKNOWLEDGEMENTS.** We thank Iain Gordon, Catharina Stroppel, Peter Webb and Michael Wemyss for helpful discussions.

## References

- [1] I. Assem, D. Simson and A. Skowroński, *Elements of the Representation Theory of Associative Algebras. Volume 1: Techniques of Representation Theory*, Cambridge University Press, 2006.
- [2] D.J. Benson, *Representations and Cohomology I: Basic Representation Theory of Finite Groups and Associative Algebras*, Cambridge University Press, 1995.

- [3] W.D. Burgess, K.R. Fuller, E.R. Voss and B. Zimmermann-Huisgen, The Cartan matrix as an indicator of finite global dimension for Artinian rings, *Proc. A.M.S.* **95** (1985) 157–165.
- [4] S. Eilenberg, Algebras of cohomologically finite dimension, *Comment. Math. Helv.* **28** (1954) 310–319.
- [5] K.R. Fuller, The Cartan determinant and global dimension of artinian rings, *Contemp. Math.* **124** (1992) 51–72.
- [6] F.W. Lawvere, Metric spaces, generalized logic and closed categories, *Rend. Sem. Mat. Fis. Milano* **XLIII** (1973) 135–166. Also *Repr. Theory Appl. Categ.* **1** (2002) 1–37.
- [7] T. Leinster, The bijection between projective indecomposable and simple modules, *Bull. Belg. Math. Soc. Simon Stevin* **22** (2015) 725–735.
- [8] T. Leinster, The Euler characteristic of a category, *Doc. Math.* **13** (2008) 21–49.
- [9] T. Leinster, The magnitude of metric spaces, *Doc. Math.* **18** (2013) 857–905.
- [10] T. Leinster and S. Willerton, On the asymptotic magnitude of subsets of Euclidean space, *Geom. Ded.* **164** (2013) 287–310.
- [11] M.W. Meckes, Positive definite metric spaces, *Positivity* **17** (2013) 733–757.
- [12] M.W. Meckes, Magnitude, diversity, capacities, and dimensions of metric spaces, *Potential Anal.* **42** (2015) 549–572.
- [13] A. Skowroński and K. Yamagata, *Frobenius Algebras I: Basic Representation Theory*, European Mathematical Society, 2012.
- [14] C.A. Weibel, *An Introduction to Homological Algebra*, Cambridge University Press, 1994.
- [15] D. Zacharia, On the Cartan matrix of an artin algebra of global dimension two, *J. Algebra* **82** (1983) 353–357.

*JC: Centre for Mathematical Science, City University  
London EC1V 0HB, United Kingdom*

*AK: Mathematical Sciences, University of Bath  
Bath BA2 7AY, United Kingdom*

*TL: School of Mathematics, University of Edinburgh  
Edinburgh EH9 3FD, United Kingdom*

Email: `j.chuang@city.ac.uk`  
`a.d.king@bath.ac.uk`  
`tom.leinster@ed.ac.uk`

This article may be accessed at <http://www.tac.mta.ca/tac/>

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.

Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

Full text of the journal is freely available from the journal's server at <http://www.tac.mta.ca/tac/>. It is archived electronically and in printed paper format.

**SUBSCRIPTION INFORMATION** Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to [tac@mta.ca](mailto:tac@mta.ca) including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, [rrosebrugh@mta.ca](mailto:rrosebrugh@mta.ca).

**INFORMATION FOR AUTHORS** The typesetting language of the journal is  $\text{\TeX}$ , and  $\text{\LaTeX}2\text{e}$  is required. Articles in PDF format may be submitted by e-mail directly to a Transmitting Editor. Please obtain detailed information on submission format and style files at <http://www.tac.mta.ca/tac/>.

**MANAGING EDITOR.** Robert Rosebrugh, Mount Allison University: [rrosebrugh@mta.ca](mailto:rrosebrugh@mta.ca)

**$\text{\TeX}$  NICAL EDITOR.** Michael Barr, McGill University: [barr@math.mcgill.ca](mailto:barr@math.mcgill.ca)

**ASSISTANT  $\text{\TeX}$  EDITOR.** Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: [gavin\\_seal@fastmail.fm](mailto:gavin_seal@fastmail.fm)

**TRANSMITTING EDITORS.**

Clemens Berger, Université de Nice-Sophia Antipolis: [cberger@math.unice.fr](mailto:cberger@math.unice.fr)

Richard Blute, Université d' Ottawa: [rblute@uottawa.ca](mailto:rblute@uottawa.ca)

Lawrence Breen, Université de Paris 13: [breen@math.univ-paris13.fr](mailto:breen@math.univ-paris13.fr)

Ronald Brown, University of North Wales: [ronnie.profbrown@btinternet.com](mailto:ronnie.profbrown@btinternet.com)

Valeria de Paiva, Nuance Communications Inc: [valeria.depaiva@gmail.com](mailto:valeria.depaiva@gmail.com)

Ezra Getzler, Northwestern University: [getzler@northwestern.edu](mailto:getzler@northwestern.edu)

Kathryn Hess, Ecole Polytechnique Fédérale de Lausanne: [kathryn.hess@epfl.ch](mailto:kathryn.hess@epfl.ch)

Martin Hyland, University of Cambridge: [M.Hyland@dpmms.cam.ac.uk](mailto:M.Hyland@dpmms.cam.ac.uk)

Anders Kock, University of Aarhus: [kock@imf.au.dk](mailto:kock@imf.au.dk)

Stephen Lack, Macquarie University: [steve.lack@mq.edu.au](mailto:steve.lack@mq.edu.au)

F. William Lawvere, State University of New York at Buffalo: [wlawvere@buffalo.edu](mailto:wlawvere@buffalo.edu)

Tom Leinster, University of Edinburgh: [Tom.Leinster@ed.ac.uk](mailto:Tom.Leinster@ed.ac.uk)

Ieke Moerdijk, Radboud University Nijmegen: [i.moerdijk@math.ru.nl](mailto:i.moerdijk@math.ru.nl)

Susan Niefield, Union College: [niefiels@union.edu](mailto:niefiels@union.edu)

Robert Paré, Dalhousie University: [pare@mathstat.dal.ca](mailto:pare@mathstat.dal.ca)

Jiri Rosicky, Masaryk University: [rosicky@math.muni.cz](mailto:rosicky@math.muni.cz)

Giuseppe Rosolini, Università di Genova: [rosolini@disi.unige.it](mailto:rosolini@disi.unige.it)

Alex Simpson, University of Ljubljana: [Alex.Simpson@fmf.uni-lj.si](mailto:Alex.Simpson@fmf.uni-lj.si)

James Stasheff, University of North Carolina: [jds@math.upenn.edu](mailto:jds@math.upenn.edu)

Ross Street, Macquarie University: [street@math.mq.edu.au](mailto:street@math.mq.edu.au)

Walter Tholen, York University: [tholen@mathstat.yorku.ca](mailto:tholen@mathstat.yorku.ca)

Myles Tierney, Université du Québec à Montréal: [tierney.myles4@gmail.com](mailto:tierney.myles4@gmail.com)

R. J. Wood, Dalhousie University: [rjwood@mathstat.dal.ca](mailto:rjwood@mathstat.dal.ca)